

## Longitudinal Study Of Peak Expiratory Flow Rate In Pregnant Women

Dr Monika Bansal\*, Dr Manoj Goyal\*\*, Dr Jasjeet Kaur Dhillon\*\*\*, Dr Parmjit Kaur\*\*\*\*

\*Assistant Professor, Dept. of Physiology & \*\*Associate Professor, Deptt. of Pharmacology, Maharishi Markandeshwar Institute of Medical Sciences & Research, Mullana, Ambala, Haryana \*\*\*Professor, Dept. of Physiology & \*\*\*\*Professor, Deptt. of Obst. & Gynae, Govt. Medical College & Rajindra Hospital, Patiala, Punjab, India

**Abstracts:** Background: Several changes have been reported in the maternal pulmonary function tests during pregnancy. A longitudinal study was undertaken to document these changes throughout pregnancy using Peak Expiratory Flow Rate (PEFR). Effect of age and height on PEFR was also documented. Method :The study included 100 pregnant females and 100 non-pregnant female controls. PEFR was measured in each trimester of pregnancy at postpartum with Mini Wright peak flow meter and the highest value of PEFR from three correctly performed blows was considered. Results: There was a decrease in mean PEFR as the pregnancy advanced from 1st to 3rd trimester and increase in PEFR in post-partum, both being statistically significant. PEFR had significant negative correlation with age. Mean PEFR increased with an increase in age of the study subjects in all the 3 trimesters of pregnancy, with maximum value at 24 -29 years of age and there after started declining. PEFR had highly significant positive correlation with height in all the 3 trimesters of pregnancy. Conclusion : The study documented the changes in PEFR during pregnancy, the effect of age and height on PEFR along with their prediction equations. [ Bansal M et al NJIRM 2012; 3(1) : 34-38]

**Key Words:** Peak Expiratory Flow Rate, Pregnancy, Active breathing exercises

**Author for correspondence:** Dr. Monika Bansal, Assistant Professor, Department of Physiology, Maharishi Markandeshwar Institute of Medical Sciences & Research, Mullana, Ambala, Haryana, India, e-mail: dr\_manojgoyal@yahoo.co.in

**Introduction:** Pregnancy is such a remarkable state of physiological adaptation, in which profound alterations in the functioning of all the systems of the mother occur to accommodate the needs of the developing fetus<sup>1</sup>. Several changes reported in the maternal pulmonary function tests during pregnancy are also a part of this adaptation<sup>2</sup>.

Increasing size of the fetus impedes the normal process of ventilation in the mother<sup>3</sup>. So, it would be logical to expect an increase in the respiratory function because the fetus depends on the mother's lungs for oxygenation and any impairment in the mother may result in fetal distress<sup>4</sup>.

Pulmonary Function Tests (PFTs), provide an accurate knowledge of the physiological changes in the pulmonary functions occurring during pregnancy, so proper evaluation of any respiratory ailment during pregnancy can be done<sup>2</sup>. Moreover, their precise knowledge allows the clinician to verify the extent of the adaptation in pregnant women and helps to avoid unnecessary treatment of physiological changes misinterpreted as pathological changes in reference to pre-pregnancy standards<sup>5</sup>. Assessment of pulmonary functions in normal women during pregnancy is also necessary

for better antenatal care, in the assessment of fitness for anesthesia and to know the progress of pre-existing lung disease<sup>6</sup>.

Peak Expiratory Flow Rate (PEFR) is an important PFT, which has been used effectively and economically by many researchers. The measurement of peak expiratory flow rate is a simple procedure in which an individual takes a full inspiration and blows out as forcibly as possible into an instrument called a peak flow meter, which measures the maximal gas flow during exhalation in litres per minute (LPM). It is also recommended by the British Thoracic Society as a useful tool in the diagnosis and management of asthma<sup>7</sup>.

Studies have been done on PEFR in pregnant women, but, with conflicting results<sup>6,8,9,10,11</sup>. So a longitudinal study was undertaken to document the changes in PEFR in each trimester of pregnancy and in postpartum and comparing them with each other and with the non-pregnant females in the same age group taken as controls. Additionally the effect of age and height on PEFR in pregnancy was also studied. Various prediction equations were also formulated.

**Material and Methods:** This longitudinal study was conducted in the department of Physiology in association with the Deptt. of Obstetrics and Gynecology at Government Medical College and Rajindra hospital, Patiala, Punjab after approval from the institutional ethics committee. The study included 100 healthy pregnant women in the age group of 18-35 years as study subjects taken from the out patient department. 100 healthy non-pregnant women in the same age group served as control taken from hospital staff and students.

The subjects were judged to be healthy on the following criteria:

- No history, current or past of any cardiovascular or respiratory disorder.
- No history of smoking.
- No history of exertional dyspnoea or general debility.
- No history of recurrent or persistent expectoration.
- No history of asthma or recurrent bronchitis during their childhood.
- No history of occupational exposure to lung toxins.
- No sign of any bony deformity of the thoracic cage.

A detailed history was taken to rule out any significant illness. A detailed general physical and obstetrical examination was done to rule out any

abnormality. Written informed consent was obtained from all subjects.

PEFR was measured with Mini Wright peak flow meter (Clement Clarke) and the highest value of PEFR from three correctly performed blows was considered. Adequate rest was given in between the readings. Before performing the procedure, it was thoroughly explained to each subject.

Data gathered was analyzed using analysis of variance (ANOVA) and 't' test. Correlation coefficient was performed by Pearson correlation analysis (r). Values of  $p < 0.05$ ,  $< 0.01$  and  $> 0.05$  were taken as statistically significant, highly significant and not significant respectively.

**Result & Discussion:** The following observations and results were drawn out:-

Table I shows the baseline characteristics of the study subjects at different trimesters and postpartum and in the controls. There was no statistically significant difference ( $p > 0.05$ ) in mean age, weight, height, body surface area (BSA) between the study subjects at postpartum and in controls. The difference in mean Hb was statistically significant in two groups.

**Table I: Baseline Characteristics Of The Study Subjects**

Variable	1 <sup>st</sup> Trimester	2 <sup>nd</sup> Trimester	3 <sup>rd</sup> Trimester	Postpartum (6-8 wks)	Control Subjects
Age (years)	25.24± 4.07	25.24± 4.07	25.24±4.07	25.24±4.07	24.92±4.62
Weight (kg)	55.26±6.12	57.75±5.62	62.13±5.74	54.81±5.22	54.31±4.06
Height (cm)	153.03± 2.93	153.03±2.93	153.03±2.93	153.03±2.93	153.28±2.96
BSA (m <sup>2</sup> )	1.514± 0.060	1.543±0.063	1.592±0.062	1.509±0.063	1.506±0.062
Hb(gm%)	10.79± 0.58	10.34±0.70	10.31±0.51	13.34±0.49	11.03±0.67

The mean PEFR of the study subjects showed statistically highly significant decline as the pregnancy advanced from 1<sup>st</sup> to 3<sup>rd</sup> trimester (Table IIb). This observation was similar to the observations by the authors in the earlier studies<sup>2,6,9,11,12,13,14,15,16,17</sup>. However, statistically highly significant increase in mean PEFR was observed at 6-8 weeks of postpartum, when compared to each trimester. This observation was consistent with the observations of the authors in the earlier studies<sup>2,13,17</sup>. However, Brancazio et al<sup>8</sup> observed not significant change during the 3 trimesters and postpartum. On the other side the

difference in mean PEFR in study subjects at postpartum and in controls was not significant statistically (Table IIb).

The variation in Mean PEFR was also found to be highly significant statistically (ANOVA) when comparison was made in between the trimesters and postpartum. (Table IIa)

The decrease in mean PEFR may be attributed to lesser force of contraction of main expiratory muscles viz. anterior abdominal muscles and internal intercostal muscles or could be due to

mechanical effect of enlarging gravid uterus affecting vertical dimension by restricting the diaphragmatic movement.

**Table IIa : Variation of PEFR in different trimesters and post-partum**

No.	Time of Observation	No. of subjects	Range of PEFR	Mean $\pm$ SD Of PEFR	P value
I	1st trimester	100	240-374	297.52 $\pm$ 32.81	<0.01
II	2 <sup>nd</sup> trimester	100	176-312	234.77 $\pm$ 34.44	
III	3 <sup>rd</sup> trimester	100	106-260	183.81 $\pm$ 33.90	
IV	postpartum	100	270-415	334.29 $\pm$ 33.15	
V	Control subjects	50	246-410	331.46 $\pm$ 40.84	

**Table IIb : Comparison of mean PEFR between different trimesters, between trimesters and postpartum, postpartum and controls**

Comparison between groups	't'	'p'	significance
I & II	13.19	<0.001	HS
I & III	24.09	<0.001	HS
I & IV	7.88	<0.001	HS
II & III	10.54	<0.001	HS
II & IV	20.81	<0.001	HS
III & IV	31.73	<0.001	HS
IV & V	0.45	>0.05	NS

The increase in PEFR at postpartum could be possibly due to regain in strength of the muscles of anterior abdominal wall leading to return of lung functions towards normal in the postpartum period. The variation in Mean PEFR according to the age was also highly significant statistically in 3 trimesters. (Table IIIa) The correlation coefficient between PEFR and age was (r -0.147) found to be significant statistically. (Table V) This observation was consistent with other studies conducted by several workers<sup>18,19,20,21,22,23,24,25,26,27,28</sup>. However, Bhargava et al<sup>29</sup> found this correlation to be not significant.

The increase in mean PEFR with age is probably due to an increase in muscular power and rapid growth

of the airway passages. The subsequent decline in mean PEFR is due to gradual decrease in muscular power because this variable is dependent upon expiratory muscle effort, lung elastic recoil and airway size, factors which are known to reduce with advancing age.

**Table IIIa: Variation of PEFR in different age groups and in different trimesters**

age group No.	Age wise	PEFR			'p' value
		1st trim.	2nd trim.	3rd trim.	
I	18-23	299.72 $\pm$ 25.22	242.59 $\pm$ 31.31	190.91 $\pm$ 29.39	<0.01
II	24-29	307.62 $\pm$ 30.89	240.82 $\pm$ 31.10	190.25 $\pm$ 30.63	
III	$\geq$ 30	276.39 $\pm$ 38.08	211.65 $\pm$ 35.91	161.17 $\pm$ 37.40	

**Table IIIb : Comparison of mean PEFR between different age groups in each trimester**

Age group	'p' value in different trimesters		
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>
I & II	>0.05	>0.05	>0.05
I & III	<0.05	<0.01	<0.01
II & III	<0.01	<0.01	<0.01

The increase in mean PEFR with the increase in height was observed in each trimester, though the mean PEFR decreased in each height group as the pregnancy advanced from 1<sup>st</sup> trimester to 3<sup>rd</sup> trimester. The increase in mean PEFR was not significant statistically on comparing height intervals I & II, but it was significant on comparing height intervals II & III and I & III (Table IVb).

**Table IVa : Variation of PEFR in different height intervals and in different trimesters**

Ht group	Ht. Int.	PEFR			'p' value
		1 <sup>st</sup> trim.	2 <sup>nd</sup> trim.	3 <sup>rd</sup> trim.	
I	145-149	275.61 $\pm$ 20.35	216.61 $\pm$ 29.44	161.84 $\pm$ 19.97	<0.01
II	150-154	291.01 $\pm$ 33.42	227.01 $\pm$ 33.35	175.82 $\pm$ 32.59	
III	$\geq$ 155	316.27 $\pm$ 28.81	254.60 $\pm$ 29.34	205.69 $\pm$ 29.10	

But, there was decrease in mean PEFR in all the height intervals as the pregnancy advanced from 1<sup>st</sup> to 3<sup>rd</sup> trimester. The decrease in mean PEFR was

highly significant statistically on comparing 1<sup>st</sup> trimester with 2<sup>nd</sup> trimester and 3<sup>rd</sup> trimester, as well as on comparing 2<sup>nd</sup> trimester with the 3<sup>rd</sup> trimester.(Table IVb)

**Table IVb : Comparison of mean PEFR between different ht. intervals in each trimester**

Ht. group	'p' value in different trimesters		
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>
I & II	>0.05	>0.05	>0.05
I & III	<0.001	<0.01	<0.01
II & III	<0.01	<0.01	<0.01

The variation in Mean PEFR according to the height was also highly significant statistically in 3 trimesters. (Table IVa)

The coefficient of correlation between PEFR and height was found to be (r +0.327) highly significant statistically.(Table V) This observation was consistent with the findings of earlier studies<sup>23,30,31,32,33,34,35,36,37,38,39,40,41</sup>. This observation could possibly be due to more chest volume in the taller subjects and an increase in expiratory muscle effort with an increase in height.

**Table V : Correlation coefficient between PEFR and age and PEFR and height**

Variable	PEFR & Age	PFR & Height
Correlation coefficient (r)	-0.147	+0.327
'p; value	<0.05	<0.001

**Table VI :Possible regression of PEFR in relation to age and height**

Variable	Regression equation
Age (years)	Y=291.175-2.07x
Height (cms)	Y=-746.46+6.43x

**Conclusion:** The present study was done to draw conclusive evidence as to what influence the normal pregnancy has on pulmonary functions. Thus, our study documented the changes in PEFR values with advancing gestational age, age and height and concluded that PEFR had significant negative correlation with age. Mean PEFR increased with an increase in age of the study subjects in all the 3 trimesters of pregnancy, with maximum value at 24-29 years of age and there after started declining. PEFR had highly significant positive correlation with height in all the 3 trimesters of pregnancy. There

was statistically highly significant decrease in mean PEFR as the pregnancy advanced from 1<sup>st</sup> to 3<sup>rd</sup> trimester and postpartum. The PEFR regressed on independent variables age and height.

#### References:

1. Das TK, Jana H. Maternal airways function during normal pregnancy. Indian J Med Sci 1991;45(10):265-8.
2. Phatak MS, Kurhade GA. A longitudinal study of antenatal changes in lung function tests and importance of postpartum exercises in their recovery. Indian J Physiol Pharmacol 2003;47(3):352-6.
3. Saxena SC, Rao VSC, Mudgal S. A study of pulmonary function tests during pregnancy. J Obstet Gynaecol India 1979;29(4):993-5.
4. Bhatia P, Bhatia K. Pregnancy and the lungs. Postgrad Med J 2000;76: 683-9.
5. Neeraj, Sodhi C, Pramod J, Singh J, kaur V. effect of advanced uncomplicated pregnancy on pulmonary function parameters of north indian subjects indian j physiol pharmacol 2010 : 54(1) : 69-72.
6. Mokkapatti R, Prasad EC, Venkatraman, Fatima K. Ventilatory functions in pregnancy. Indian J Physiol Pharmacol 1991;35(4):237-40.
7. Higgins D, Measuring PEFR. Nursing Practice,clinical research. 2005 March. 101(10): 32
8. Brancazio LR, Laifer SA, Schwartz T. Peak expiratory flow rate in normal pregnancy. Obstet Gynecol 1997;89(3):383-6.
9. Monga U, Kumari K. Pulmonary functions in Punjabi pregnant women. Indian J Physiol Pharmacol 2000;44(1):115-6.
10. Rubin A, Russo N, Goucher D. The effect of pregnancy upon pulmonary function in normal women. Am J Obstet Gynaecol 1956; 72(5):963-9.
11. Ganerwal SK, Deshpande DR, Reddy BV, Shaikh RM. Effect of pregnancy on pulmonary ventilation. J Obstet Gynaecol India 1984;36(3):639-41.
12. Singhal U, Saxena K. Effect of anaemia on respiratory and metabolic parameters during third trimester of pregnancy. Indian J Physiol Pharmacol 1987;31(2):130-5.
13. Puranik BM, Kurhade GA, Kaore SB, Patwardhan SA, Kher JR. PEFR in pregnancy: a longitudinal

- study. *Indian J Physiol Pharmacol* 1995;39(2):135-9.
14. Rasheed BMA, Hussain K, Hussain S. PEFR in relation to phases of pregnancy. *Indian J Physiol Pharmacol* 2000;44(4):511-2.
  15. Rasheed BMA, Mansoor A, Hussain S. Incentive spirometry and PEFR in different phases of pregnancy. *Indian J Physiol Pharmacol* 2002;46(1):126-8.
  16. Sroczyński T. Evaluation of respiratory tract function in healthy women in the last month of uncomplicated pregnancy. *Ann Acad Med Stetin* 2002;48:331-50.
  17. Harirah HM, Donia SE, Nasrallah FK, Saade GR, Belfort MA. Effect of gestational age and position on peak expiratory flow rate: a longitudinal study. *Obstet Gynaecol* 2005;105(2):372-6.
  18. Flint FJ, Khan MO. Clinical use of peak flow meter. *Br Med J* 1962;2:1231-2.
  19. Pelzer AM, Thomson ML. Expiratory peak flow. *Br Med J* 1964;2:123.
  20. Gupta S, Puri MB, Singh SI. Pulmonary function tests in health. *J Assoc Physiol India* 1975;23:247-52.
  21. Cookson JB, Blake GTW, Faranisi C. Normal values for ventilatory function in Rhodesian Africans. *Br J Dis Chest* 1976;70:107-11.
  22. Gupta P, Gupta S, Ajmera RL. Lung function tests in Rajasthani subjects. *Indian J Physiol Pharmacol* 1979;23(1):8-14.
  23. Singh HD, Peri S. PEFR in south Indian adults. *Indian J Physiol Pharmacol* 1979;23(4):315-20.
  24. Gupta CK, Mathur N. Statistical models relating PEFR to age, height and weight in men and women. *J Epidemiol Commun Hlth* 1982;36:64-7.
  25. Jain SK, Kumar R, Sharma DA. Factors influencing PEFR in normal subjects. *Lung India* 1983;3:92-7.
  26. Lebowitz MD, Sherrill DL, Kaltenborn W, Burrows B. Peak expiratory flow from maximum expiratory flow volume curves in a community population :cross- sectional and longitudinal analyses. *Eur Respir J Suppl* 1997;24:29s-38s.
  27. Nishitsuji M, Fujimura M, Shibata K. Longitudinal decline of forced expiratory volume in one second in non-smoking Japanese women. *Nihon Kokyuki Gakkai Zasshi* 2006;44(4):301-4
  28. Prasad R, Verma SK, Agrawal GG, Mathur N. Prediction model for peak expiratory flow in north Indian population. *Indian J Chest Dis Allied Sci.* 2006;48(2):103-6.
  29. Bhargava RP, Misra SM, Gupta NK. Ventilatory tests and lung volume studies in Madhya Pradesh : physiological norms. *Indian J Physiol Pharmacol* 1973;17(3):267-72.
  30. Leiner GC, Abramowitz S, Small MJ, Stenby VB, Lewis WA. Peak expiratory flow rate. Standard values for normal subjects. Use as a clinical test of ventilator function. *Am Rev Respir Dis* 1963;88: 644-51.
  31. Elebute EA, Femi-Pearse D. Peak flow rate in Nigeria : anthropometric determinants and usefulness in assessment of ventilatory function. *Thorax* 1971;26:597- 601.
  32. Woolcock AJ, Colman MH, Blackburn CRB. Factors affecting normal values for ventilatory lung functions. *Am Rev Respir Dis* 1972;106:692-709.
  33. Amin SK, Pande RS. PEFR in normal subjects. *India J Chest Dis.* 1978;20:81-3.
  34. Bayu T, Teshale S, Mills RJ. PEFR in normal Ethiopian children and adults in Addis Ababa. *Br J Dis Chest* 1987; 81:176.
  35. Udawadia FE, Sunavala JD, Shetye VM. Lung function studies in healthy Indian subjects. *J Appl Physiol India* 1987;36(7):491-6.
  36. Mukhtar MSR, Rao GMM, Morghom LO. Peak expiratory flow rates in Libyan adolescents. *Indian J Physiol Pharmacol* 1989;33(4):223-7.
  37. Walter S, Richard J. Lung function development in Indian men and women during late adolescence and early adulthood : a longitudinal study. *Indian J Physiol Pharmacol* 1991;35(1):15-20.
  38. Rao NM, Mavlankar MG, Kulkarni PK, Kashyap SK. Pulmonary function studies in Gujrati subjects. *Indian J Physiol Pharmacol* 1992;36(1):55-9.
  39. Jepegnanam V, Amirtharaj G, Sulochana, Damodarasamy, Rao VM. PEFR in a random healthy population of Coimbatore. *India J Physiol Pharmacol* 1996;40(2):127-33.
  40. Mohamed El, Maiolo C, Iacopino L, Pepe M, Di Daniele N, De Lorenzo A. The impact of body-weight components on forced spirometry in healthy Italians. *Lung* 2002;180(3):149-59.
  41. Dikshit MB, Raje S, Agrawal MJ. Lung functions with spirometry: an Indian perspective-I. Peak expiratory flow rates. *Indian J Physiol Pharmacol* 2005;49(1):8-18.